arXiv:1001.0034v1 [math.NT] 4 Jan 2010

NEW IDENTITIES INVOLVING q-EULER POLYNOMIALS OF HIGHER ORDER

T. Kim AND Y. H. Kim

Abstract. In this paper, we present new generating functions which are related to q-Euler numbers and polynomials of higher order. From these generating functions, we give new identities involving q-Euler numbers and polynomials of higher order.

§1. Introduction/ Preliminaries

Let $\mathbb{C}$ be the complex number field. We assume that $q \in \mathbb{C}$ with $|q| < 1$ and the q-number is defined by $[x]_q = \frac{1-q^x}{1-q}$ in this paper. The q-factorial is given by $[n]_q! = [n]_q[n-1]_q \cdots [2]_q[1]_q$ and the q-binomial formulae are known that

$$(x:q)_n = \prod_{i=1}^{n}(1-xq^{i-1}) = \sum_{i=0}^{n} \binom{n}{i}_q q^{\binom{i}{2}}(-x)^i, \text{ (see [3, 14, 15]),}$$

and

$$\frac{1}{(x:q)_n} = \prod_{i=1}^{n}\left(\frac{1}{1-xq^{i-1}}\right) = \sum_{i=0}^{\infty}\binom{n+i-1}{i}_q x^i, \text{ (see [3, 5, 14, 15]),}$$

where $\binom{n}{i}_q = \frac{[n]_q!}{[n-i]_q![i]_q!} = \frac{[n]_q[n-1]_q\cdots[n-i+1]_q}{[i]_q!}$.

The Euler polynomials are defined by $\frac{2}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!}$, for $|t| < \pi$. In the special case $x = 0$, $E_n(= E_n(0))$ are called the n-th Euler numbers. In this paper, we consider the q-extensions of Euler numbers and polynomials of higher order. Barnes' multiple Bernoulli polynomials are also defined by

(1)
$$\frac{t^r}{\prod_{j=1}^{r}(e^{a_jt}-1)}e^{xt} = \sum_{n=0}^{\infty} B_n(x, r|a_1, \cdots, a_r)\frac{t^n}{n!}, \text{ where } |t| < \max_{1\le i\le r}\frac{2\pi}{|a_i|}, \text{ (see [1, 14]).}$$

Key words and phrases. : multiple q-zeta function, q-Euler numbers and polynomials, higher order q-Euler numbers, Laurent series, Cauchy integral.

2000 AMS Subject Classification: 11B68, 11S80

The present Research has been conducted by the research Grant of Kwangwoon University in 2010

Typeset by $\mathcal{AMS}$-TEX

In one of an impressive series of papers (see [1, 6, 14]), Barnes developed the so-called multiple zeta and multiple gamma function. Let $a_1, \cdots, a_N$ be positive parameters. Then Barnes' multiple zeta function is defined by

$$\zeta_N(s, w|a_1, \cdots, a_N) = \sum_{m_1, \cdots, m_N=0} (w + m_1a_1 + \cdots + m_Na_N)^{-s}, \text{ (see [1])},$$

where $\Re(s) > N$, $\Re(w) > 0$. For $m \in \mathbb{Z}_+$, we have

$$\zeta_N(-m, w|a_1, \cdots, a_N) = \frac{(-1)^m m!}{(N+m)!} B_{N+m}(w, N|a_1, \cdots, a_N).$$

In this paper, we consider Barnes' type multiple q-Euler numbers and polynomials. The purpose of this paper is to present new generating functions which are related to q-Euler numbers and polynomials of higher order. From the Mellin transformation of these generating functions, we derive the q-extensions of Barnes' type multiple zeta functions, which interpolate the q-Euler polynomials of higher order at negative integer. Finally, we give new identities involving q-Euler numbers and polynomials of higher order.

§2. q-Euler numbers and polynomials of higher order

In this section, we assume that $q \in \mathbb{C}$ with $|q| < 1$. Let $x, a_1, \ldots, a_r$ be complex numbers with positive real parts. Barnes' type multiple Euler polynomials are defined by

$$(2) \qquad \frac{2^r}{\prod_{j=1}^r (e^{a_jt}+1)} e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x|a_1, \ldots, a_r)\frac{t^n}{n!}, \text{ for } |t| < \max_{1\le i\le r} \frac{\pi}{|w_i|}, \text{ (see [6])},$$

and $E_n^{(r)}(a_1, \ldots, a_r)(= E_n^{(r)}(0|a_1, \ldots, a_r))$ are called the n-th Barnes' type multiple Euler numbers. First, we consider the q-extension of Euler polynomials. The q-Euler polynomials are defined by

$$(3) \qquad F_q(t, x) = \sum_{n=0}^{\infty} E_{n,q}(x)\frac{t^n}{n!} = [2]_q \sum_{m=0}^{\infty} (-q)^m e^{[m+x]_q t}, \text{ (see [8, 11, 13, 14, 15])}.$$

From (3), we have

$$E_{n,q}(x) = \frac{[2]_q}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} \frac{(-1)^l q^{lx}}{(1+q^{l+1})}.$$

In the special case $x = 0$, $E_{n,q}(= E_{n,q}(0))$ are called the n-th q-Euler numbers. From (3), we can easily derive the following relation.

$$E_{0,q} = 1, \text{ and } q(qE+1)^n + E_{n,q} = 0 \text{ if } n \ge 1, \text{ (see [8, 16, 17])},$$

where we use the standard convention about replacing E^k by $E_{k,q}$. It is easy to show that

$$\lim_{q\to 1} F_q(t,x) = \frac{2}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!}, \text{ (see [2, 3, 19-23])},$$

where $E_n(x)$ are the n-th Euler polynomials. For $r \in \mathbb{N}$, the Euler polynomials of order r is defined by

$$\left(\frac{2}{e^t+1}\right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x)\frac{t^n}{n!}, \text{ for } |t| < \pi. \tag{4}$$

Now we consider the q-extension of (4).

$$F_q^{(r)}(t,x) = [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty} (-q)^{m_1+\cdots+m_r} e^{[m_1+\cdots+m_r+x]_q t} = \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x)\frac{t^n}{n!}, \tag{5}$$

where $E_{n,q}^{(r)}(x)$ are called the n-th q-Euler polynomials of order r (see [10-15]). From (5), we can derive

$$E_{n,q}^{(r)}(x) = \frac{[2]_q^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}\frac{(-1)^l q^{lx}}{(1+q^{l+1})^r}. \tag{6}$$

By (5) and (6), we see that

$$F_q^{(r)}(t,x) = [2]_q^r \sum_{m=0}^{\infty}\binom{m+r-1}{m}(-q)^m e^{[m+x]_q t}. \tag{7}$$

Thus, we note that $\lim_{q\to 1} F_q^{(r)}(t,x) = \left(\frac{2}{e^t+1}\right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x)\frac{t^n}{n!}$. In the special case $x = 0$, $E_{n,q}^{(r)}(= E_{n,q}^{(r)}(0))$ are called the n-th q-Euler numbers of order r. By (5), (6) and (7), we obtain the following proposition.

Proposition 1. *For $r \in \mathbb{N}$, let*

$$F_q^{(r)}(t,x) = [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty} (-q)^{m_1+\cdots+m_r} e^{[m_1+\cdots+m_r+x]_q t} = \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x)\frac{t^n}{n!}.$$

Then we have

$$E_{n,q}^{(r)}(x) = \frac{[2]_q^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}\frac{(-1)^l q^{lx}}{(1+q^{l+1})^r} = [2]_q^r \sum_{m=0}^{\infty}\binom{m+r-1}{m}(-q)^m [m+x]_q^n.$$

From the Mellin transformation of $F_q^{(r)}(t,x)$, we can derive the following equation.

$$\begin{aligned}\frac{1}{\Gamma(s)}\int_0^\infty F_q^{(r)}(-t,x)t^{s-1}dt &= [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty} \frac{(-q)^{m_1+\cdots+m_r}}{[m_1+\cdots+m_r+x]_q^s} \\ &= [2]_q^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}(-q)^m \frac{1}{[m+x]_q^s},\end{aligned} \tag{8}$$

where $s\in\mathbb{C}$, $x\neq 0,-1,-2,\dots$. By (8), we can define the multiple q-zeta function related to q-Euler polynomials.

Definition 2. *For $s\in\mathbb{C}$, $x\in\mathbb{R}$ with $x\neq 0,-1,-2,\dots$, we define the multiple q-zeta function related to q-Euler polynomials as*

$$\zeta_{q,r}(s,x) = [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty} \frac{(-q)^{m_1+\cdots+m_r}}{[m_1+\cdots+m_r+x]_q^s}.$$

Note that $\zeta_{q,r}(s,x)$ is a meromorphic function in whole complex s-plane. From (8), we also note that

$$\zeta_{q,r}(s,x) = [2]_q^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}(-q)^m \frac{1}{[m+x]_q^s}.$$

By Laurent series and the Cauchy residue theorem in (5) and (8), we see that

$$\zeta_q(-n,x) = E_{n,q}^{(n)}(x), \text{ for } n\in\mathbb{Z}_+.$$

Therefore, we obtain the following theorem.

Theorem 3. *For $r\in\mathbb{N}, n\in\mathbb{Z}_+$, and $x\in\mathbb{R}$ with $x\neq 0,-1,-2,\dots$, we have*

$$\zeta_q(-n,x) = E_{n,q}^{(r)}(x).$$

Let χ be the Dirichlet's character with conductor $f\in\mathbb{N}$ with $f\equiv 1 \pmod 2$. Then the generalized q-Euler polynomial attached to χ are considered by

$$F_{q,\chi}(x) = \sum_{n=0}^{\infty} E_{n,\chi,q}(x)\frac{t^n}{n!} = [2]_q \sum_{m=0}^{\infty}(-q)^m\chi(m)e^{[m+x]_q t}.$$

From (3) and (9), we have

$$E_{n,\chi,q}(x) = \frac{[2]_q}{[2]_{q^f}} \sum_{a=0}^{f-1}(-q)^a\chi(a)E_{n,q^f}(\frac{x+a}{f}).$$

In the special case $x = 0$, $E_{n,\chi,q} = E_{n,\chi,q}(0)$ are called the n-th generated q-Euler number attached to χ.

It is known that the generalized Euler polynomials of order r are defined by

$$(\frac{2\sum_{a=0}^{f-1}(-1)^a\chi(a)e^{at}}{e^{ft}+1})^r e^{xt} = \sum_{n=0}^{\infty} E_{n,\chi}^{(r)}(x)\frac{t^n}{n!}, \tag{10}$$

for $|t| < \frac{\pi}{f}$.

We consider the q-extension of (10). The generalized q-Euler polynomials of order r attached to χ are defined by

$$\begin{aligned} F_{q,\chi}^{(r)}(t,x) &= [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty} (-q)^{m_1+\dots+m_r}(\prod_{i=1}^{r}\chi(m_i))e^{[m_1+\dots+m_r+x]_q t} \\ &= \sum_{n=0}^{\infty} E_{n,\chi,q}^{(r)}(x)\frac{t^n}{n!}, \text{ (see [14, 15]).} \end{aligned} \tag{11}$$

Note that

$$\lim_{q\to 1} F_{q,\chi}^{(r)}(t,x) = (\frac{2\sum_{a=0}^{f-1}(-1)^a\chi(a)e^{at}}{e^{ft}+1})^r.$$

By (11), we easily see that

$$\begin{aligned} E_{n,\chi,q}^{(r)}(x) &= \frac{[2]_q^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}(-q^x)^l \sum_{a_1,\dots,a_r=0}^{f-1}(\prod_{j=1}^{r}\chi(a_j))\frac{(-q^{l+1})^{\sum_{i=1}^r a_i}}{(1+q^{(l+1)f})^r} \\ &= [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty}(-q)^{m_1+\dots+m_r}(\prod_{i=1}^{r}\chi(m_i))[m_1+\dots+m_r+x]_q^n. \end{aligned}$$

For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$, we have

$$\begin{aligned} &\frac{1}{\Gamma(s)}\int_0^{\infty} F_{q,\chi}^{(r)}(-t,x)t^{s-1}dt \\ &= [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty}\frac{(-q)^{m_1+\dots+m_r}(\prod_{i=1}^{r}\chi(m_i))}{[m_1+\dots+m_r+x]_q^s}, \text{ (see [15]).} \end{aligned} \tag{12}$$

From (12), we can consider the Dirichlet's type multiple q-l-function as follows :

Definition 4. *For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots,$ we define the Dirichlet's type multiple q-l-function as*

$$l_q(s,x|\chi) = [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty}\frac{(-q)^{m_1+\dots+m_r}(\prod_{i=1}^{r}\chi(m_i))}{[m_1+\dots+m_r+x]_q^s}, \text{ (see [15]).}$$

By Laurent series and the Cauchy residue theorem in (11) and (12), we obtain the following theorem.

Theorem 5. *For $n \in \mathbb{Z}_+$, we have*

$$l_q(-n, x|\chi) = E^{(r)}_{n,\chi,q}(x).$$

For $h \in \mathbb{Z}$ and $r \in \mathbb{N}$, we consider the extended r-ple q-Euler polynomials.

$$\begin{aligned} F_q^{(h,r)}(t,x) &= [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty} q^{\sum_{j=1}^r (h-j+1)m_j}(-1)^{\sum_{j=1}^r m_j} e^{[m_1+\cdots+m_r+x]_q t} \\ &= \sum_{n=0}^{\infty} E_{n,q}^{(h,r)}(x)\frac{t^n}{n!}. \end{aligned} \tag{13}$$

Note that

$$\lim_{q\to 1} F_q^{(h,r)}(t,x) = (\frac{2}{e^t+1})^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x)\frac{t^n}{n!}.$$

From (13), we note that

$$\begin{aligned} E_{n,q}^{(h,r)}(x) &= \frac{[2]_q^r}{(1-q)^n}\sum_{l=0}^{n}\binom{n}{l}\frac{(-q^x)^l}{(-q^{h-r+l+1}:q)_r} \\ &= [2]_q^r \sum_{m=0}^{\infty}\binom{m+r-1}{m}_q (-q^{h-r+1})^m [m+x]_q^n. \end{aligned} \tag{14}$$

By (14), we easily see that

$$F_q^{(h,r)}(t,x) = [2]_q^r \sum_{m=0}^{\infty}\binom{m+r-1}{m}_q (-q^{h-r+1})^m e^{[m+x]_q t}, \text{ (see [11, 13, 14])}. \tag{15}$$

Using the Mellin transform for $F_q^{(h,r)}(t,x)$, we have

$$\begin{aligned} &\frac{1}{\Gamma(s)}\int_0^{\infty} F_q^{(r)}(-t,x)t^{s-1}dt \\ &= [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty} \frac{(-1)^{m_1+\cdots+m_r} q^{\sum_{j=1}^r (h-j+1)m_j}}{[m_1+\cdots+m_r+x]_q^s}, \text{ (see [13, 14, 15])}, \end{aligned} \tag{16}$$

for $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \ldots$. Now we can define the extended q-zeta function associated with $E_{n,q}^{(h,r)}(x)$.

Definition 6. *For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \ldots,$ we define the (h, q)-zeta function as*

$$\zeta_{q,r}^{(h)}(s,x) = [2]_q^r \sum_{m_1,\ldots,m_r=0}^{\infty} \frac{(-1)^{m_1+\cdots+m_r} q^{\sum_{j=1}^r (h-j+1)m_j}}{[m_1+\cdots+m_r+x]_q^s}.$$

Note that $\zeta_{q,r}^{(h)}(s,x)$ is also a meromorphic function in whole complex s-plane. From (16) and (15), we note that

$$\zeta_{q,r}^{(h)}(s,x) = [2]_q^r \sum_{m=0}^{\infty} \binom{m+r-1}{m}_q (-q^{h-j+1})^m \frac{1}{[m+x]_q^s}. \tag{17}$$

Using the Cauchy residue theorem and Laurent series in (16), we obtain the following theorem.

Theorem 7. *For $n \in \mathbb{Z}_+$, we have*

$$\zeta_{q,r}^{(h)}(-n,x) = E_{n,q}^{(h,r)}(x).$$

We consider the extended r-ple generalized q-Euler polynomials as follows :

$$\begin{aligned} &F_{q,\chi}^{(h,r)}(t,x) \\ &= [2]_q^r \sum_{m_1,\ldots,m_r=0}^{\infty} q^{\sum_{j=1}^r (h-j+1)m_j} (-1)^{\sum_{j=1}^r m_j} (\prod_{j=1}^r \chi(m_j)) e^{[m_1+\cdots+m_r+x]_q t} \\ &= \sum_{n=0}^{\infty} E_{n,\chi,q}^{(h,r)}(x) \frac{t^n}{n!}. \end{aligned} \tag{18}$$

By (18), we see that

$$\begin{aligned} E_{n,\chi,q}^{(h,r)}(x) &= \frac{[2]_q^r}{(1-q)^n} \sum_{a_1,\ldots,a_r=0}^{f-1} (-1)^{\sum_{j=1}^r a_j} (\prod_{j=1}^r \chi(a_j)) \sum_{l=0}^{n} \binom{n}{l} \frac{(-1)^l q^{lx} q^{(h-j+l+1)a_j}}{(-q^{(h-r+l+1)f} : q^f)_r} \\ &= \frac{[2]_q^r}{[2]_{q^f}^r} [f]_q^n \sum_{a_1,\ldots,a_r=0}^{f-1} (-1)^{\sum_{j=1}^r a_j} (\prod_{j=1}^r \chi(a_j)) q^{\sum_{j=1}^r (h-j+1)a_j} \zeta_{q^f,r}^{(h)}(-n, \frac{x+\sum_{j=1}^r a_j}{f}). \end{aligned} \tag{19}$$

Therefore, we obtain the following theorem.

Theorem 8. *For $n \in \mathbb{Z}_+$, we have*

$$E_{n,\chi,q}^{(h,r)}(x) \\ = \frac{[2]_q^r}{[2]_{q^f}^r}[f]_q^n \sum_{a_1,\dots,a_r=0}^{f-1} (-1)^{\sum_{j=1}^r a_j} (\prod_{j=1}^r \chi(a_j)) q^{\sum_{j=1}^r (h-j+1)a_j} \zeta_{q^f,r}^{(h)}(-n, \frac{x+\sum_{j=1}^r a_j}{f}).$$

From (18), we note that

$$\frac{1}{\Gamma(s)}\int_0^\infty F_{q,\chi}^{(h,r)}(-t,x)t^{s-1}dt \\ = [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty} \frac{q^{\sum_{j=1}^r (h-j+1)m_j}(\prod_{j=1}^r \chi(m_j))(-1)^{m_1+\cdots+m_r}}{[m_1+\cdots+m_r+x]_q^s}, \tag{20}$$

where $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$.

From (20), we define the Dirichlet's type multiple (h,q)-l-function associated with the generalized multiple q-Euler polynomials attached to χ.

Definition 9. *For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$, we define the Dirichlet's type multiple q-l-function as follows :*

$$l_q^{(h)}(s,x|\chi) = [2]_q^r \sum_{m_1,\dots,m_r=0}^{\infty} \frac{q^{\sum_{j=1}^r (h-j+1)m_j}(\prod_{i=1}^r \chi(m_i))(-1)^{m_1+\cdots+m_r}}{[m_1+\cdots+m_r+x]_q^s}.$$

Note that $l_q^{(h)}(s,x|\chi)$ is a meromorphic function in whole complex plane. It is easy to show that

$$l_q^{(h)}(s,x|\chi) \\ = \frac{[2]_q^r}{[2]_{q^f}^r}\frac{1}{[f]_q^s} \sum_{a_1,\dots,a_r=0}^{f-1} (-1)^{\sum_{j=1}^r a_j} (\prod_{j=1}^r \chi(a_j)) q^{\sum_{j=1}^r (h-j+1)a_j} \zeta_{q^f,r}^{(h)}(s, \frac{x+\sum_{j=1}^r a_j}{f}).$$

By (19) and (20), we obtain the following theorem.

Theorem 10. *For $n \in \mathbb{Z}_+$, we have*

$$l_q^{(h)}(-n,x|\chi) = E_{n,\chi,q}^{(h,r)}(x).$$

Finally, we give the q-extension of Barnes' type multiple Euler polynomials in (2). For $x, a_1, \dots, a_r \in \mathbb{C}$ with positive real part, let us define the Barnes' type mutiple

q-Euler polynomials in $\mathbb{C}$ as follows :

$$\begin{aligned} & F_q^{(r)}(t, x|a_1, \dots, a_r; b_1, \dots, b_r) \\ & = [2]_q^r \sum_{m_1, \dots, m_r = 0}^{\infty} (-1)^{m_1 + \cdots + m_r} q^{(b_1+1)m_1 + \cdots + (b_r+1)m_r} e^{[a_1 m_1 + \cdots + a_r m_r + x]t} \\ & = \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x|a_1, \dots, a_r; b_1, \dots, b_r) \frac{t^n}{n!}, \end{aligned} \tag{21}$$

where $b_1, \dots, b_r \in \mathbb{Z}$. By (21), we see that

$$\begin{aligned} & E_{n,q}^{(r)}(x|a_1, \dots, a_r; b_1, \dots, b_r) \\ & = \frac{[2]_q^r}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} \frac{(-1)^l q^{lx}}{(1+q^{la_1+b_1+1}) \cdots (1+q^{la_r+b_r+1})} \\ & = [2]_q^r \sum_{m_1, \dots, m_r = 0}^{\infty} (-1)^{m_1 + \cdots + m_r} q^{(b_1+1)m_1 + \cdots + (b_r+1)m_r} [a_1 m_1 + \cdots + a_r m_r + x]_q^n. \end{aligned}$$

From (21), we note that

$$\begin{aligned} & \frac{1}{\Gamma(s)} \int_0^{\infty} F_q^{(r)}(-t, x|a_1, \dots, a_r; b_1, \dots, b_r) t^{s-1} dt \\ & = [2]_q^r \sum_{m_1, \dots, m_r = 0}^{\infty} \frac{(-q)^{m_1 + \cdots + m_r} q^{b_1 m_1 + \cdots + b_r m_r}}{[a_1 m_1 + \cdots + a_r m_r + x]_q^s}. \end{aligned} \tag{22}$$

By (22), we define the Barnes' type multiple q-zeta function as follows :

$$\begin{aligned} & \zeta_{q,r}(s, x|a_1, \dots, a_r; b_1, \dots, b_r) \\ & = [2]_q^r \sum_{m_1, \dots, m_r = 0}^{\infty} \frac{(-q)^{m_1 + \cdots + m_r} q^{b_1 m_1 + \cdots + b_r m_r}}{[a_1 m_1 + \cdots + a_r m_r + x]_q^s}, \end{aligned}$$

where $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \dots$. By (21), (22) and (23), we obtain the following theorem.

Theorem 11. *For $n \in \mathbb{Z}_+$, we have*

$$\zeta_{q,r}(s, x|a_1, \dots, a_r; b_1, \dots, b_r) = E_{n,q}^{(r)}(x|a_1, \dots, a_r; b_1, \dots, b_r).$$

Let χ be the Dirichlet's character with conductor $f \in \mathbb{N}$ with $f \equiv 1 \pmod 2$. Then the generalized Barnes' type multiple q-Euler polynomials attached to χ are defined by

$$
\begin{aligned}
& F_{q,\chi}^{(r)}(t,x|a_1,\ldots,a_r;b_1,\ldots,b_r) \\
(24) \qquad & = [2]_q^r \sum_{m_1,\ldots,m_r=0}^{\infty} (-q)^{m_1+\cdots+m_r} q^{b_1m_1+\cdots+b_rm_r} (\prod_{i=1}^{r} \chi(m_i)) e^{[a_1m_1+\cdots+a_rm_r+x]_q t} \\
& = \sum_{n=0}^{\infty} E_{n,\chi,q}^{(r)}(x|a_1,\ldots,a_r;b_1,\ldots,b_r)\frac{t^n}{n!},
\end{aligned}
$$

From (24), we note that

$$
\begin{aligned}
& \frac{1}{\Gamma(s)}\int_0^{\infty} F_{q,\chi}^{(r)}(-t,x|a_1,\ldots,a_r;b_1,\ldots,b_r)t^{s-1}dt \\
(25) \qquad & = [2]_q^r \sum_{m_1,\ldots,m_r=0}^{\infty} \frac{(-q)^{m_1+\cdots+m_r} q^{b_1m_1+\cdots+b_rm_r} (\prod_{i=1}^{r} \chi(m_i))}{[a_1m_1+\cdots+a_rm_r+x]_q^s}.
\end{aligned}
$$

By (25), we can define Barnes' type multiple q-l-function in $\mathbb{C}$. For $s \in \mathbb{C}$, $x \in \mathbb{R}$ with $x \neq 0, -1, -2, \ldots$, let us define the Barnes' type multiple q-l-function as follows :

$$
\begin{aligned}
& l_q^{(r)}(s,x|a_1,\ldots,a_r;b_1,\ldots,b_r) \\
(26) \qquad & = [2]_q^r \sum_{m_1,\ldots,m_r=0}^{\infty} \frac{(-q)^{m_1+\cdots+m_r} q^{b_1m_1+\cdots+b_rm_r} (\prod_{i=1}^{r} \chi(m_i))}{[a_1m_1+\cdots+a_rm_r+x]_q^s}.
\end{aligned}
$$

Note that $l_q^{(r)}(s,x|a_1,\ldots,a_r;b_1,\ldots,b_r)$ is a meromorphic function in whole complex s-plane. By (24), (25) and (26), we easily see that

$$
l_q^{(r)}(-n,x|a_1,\ldots,a_r;b_1,\ldots,b_r) = E_{n,\chi,q}^{(r)}(x|a_1,\ldots,a_r;b_1,\ldots,b_r)
$$

for $n \in \mathbb{Z}_+$, (see [1-18]).

References

[1] E. W. Barnes, *On the theory of multiple gamma function*, Trans. Camb. Ohilos. Soc. A **196** (1904), 374-425.

[2] I. N. Cangul,V. Kurt, H. Ozden, Y. Simsek, *On the higher-order w-q-Genocchi numbers*, Adv. Stud. Contemp. Math. **19** (2009), 39–57.

[3] N. K.Govil, V. Gupta, *Convergence of q-Meyer-Konig-Zeller-Durrmeyer operators*, Adv. Stud. Contemp. Math. **19** (2009), 97–108.

[4] T. Kim, *On a q-analogue of the p-adic log gamma functions and related integrals*, J. Number Theory **76** (1999), 320–329.

[5] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288–299.

[6] T. Kim, *On Euler-Barnes multiple zeta functions*, Russ. J. Math. Phys. **10** (2003), 261–267.

[7] T. Kim, *Analytic continuation of multiple q-zeta functions and their values at negative integers*, Russ. J. Math. Phys. **11** (2004), 71–76.
[8] T. Kim, *The modified q-Euler numbers and polynomials*, Adv. Stud. Contemp. Math. **16** (2008), 161–170.
[9] T. Kim, *Note on the q-Euler numbers of higher order*, Adv. Stud. Contemp. Math. **19** (2009), 25–29.
[10] T. Kim, *Note on Dedekind type DC sums*, Adv. Stud. Contemp. Math. **18** (2009), 249–260.
[11] T. Kim, *Note on the Euler q-zeta functions*, J. Number Theory **129** (2009), 1798–1804.
[12] T. Kim, *A note on the generalized q-Euler numbers*, Proc. Jangjeon Math. Soc. **12** (2009), 45–50.
[13] T. Kim, *Some identities on the q-Euler polynomials of higher order and q-stirling numbers by the fermionic p-adic integral on* $\mathbb{Z}_p$, Russ. J. Math. Phys. **16** (2009), 1061-9208.
[14] T. Kim, *Barnes type multiple q-zeta functions and q-Euler polynomials*, arXiv:0912.5119v1.
[15] T. Kim, *Note on multiple q-zeta functions*, to be appeared in Russ. J. Math. Phys., arXiv:0912.5477v1.
[16] T. Kim, *On the q-extension of Euler and Genocchi numbers*, J. Math. Anal. Appl. **326**, 1458–1465.
[17] T. Kim, *On p-adic q-l-functions and sums of powers*, J. Math. Anal. Appl. **329**, 1472–1481.
[18] T. Kim, Y. Simsek, *Analytic continuation of the multiple Daehee q-l-functions associated with Daehee numbers*, Russ. J. Math. Phys. **15** (2008), 58–65.
[19] Y. H. Kim, W. Kim, C. S. Ryoo, *On the twisted q-Euler zeta function associated with twisted q-Euler numbers*, Proc. Jangjeon Math. Soc. **12** (2009), 93-100.
[20] H. Ozden, I. N. Cangul, Y. Simsek, *Remarks on q-Bernoulli numbers associated with Daehee numbers*, Adv. Stud. Contemp. Math. **18** (2009), 41-48.
[21] K. Shiratani, S. Yamamoto, *On a p-adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci., Kyushu University Ser. A **39** (1985), 113-125.
[22] Y. Simsek, *Theorems on twisted L-function and twisted Bernoulli numbers*, Advan. Stud. Contemp. Math. **11** (2005), 205–218.
[23] Z. Zhang, Y. Zhang, *Summation formulas of q-series by modified Abel's lemma*, Adv. Stud. Contemp. Math. **17** (2008), 119–129.

Taekyun Kim
Division of General Education-Mathematics, Kwangwoon University,
Seoul 139-701, S. Korea e-mail: tkkim@kw.ac.kr

Young-Hee Kim
Division of General Education-Mathematics,
Kwangwoon University,
Seoul 139-701, S. Korea e-mail: yhkim@kw.ac.kr